

## University of Groningen

### SDD

Benus, R. F.; Harmsen, H. J.; van der Werf, T. S.

*Published in:*  
Intensive Care Medicine

*DOI:*  
[10.1007/s00134-010-1947-9](https://doi.org/10.1007/s00134-010-1947-9)

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2010

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

Benus, R. F., Harmsen, H. J., & van der Werf, T. S. (2010). SDD: don't be selective in considering pros and cons. *Intensive Care Medicine*, 36(10), 1785-1786. <https://doi.org/10.1007/s00134-010-1947-9>

#### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

#### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*

R. F. Benus  
H. J. Harmsen  
T. S. van der Werf

## SDD: don't be selective in considering pros and cons

Accepted: 2 May 2010  
Published online: 7 July 2010  
© The Author(s) 2010. This article is  
published with open access at  
Springerlink.com

Dear Editor,  
We thank Dr. Zandstra and  
colleagues [1] for their interest in  
our work [2]. Although in their sci-  
entific careers they have amassed  
impressive data on selective digestive  
microbial decontamination (SDD),  
they have not fully appreciated what  
our paper adds to this large body of  
knowledge.

The intestinal microbiota contains  
many more cells than the human body  
itself. Furthermore the intestinal  
microbial gene set is approximately  
150 times larger than the human  
genome [3]. Last year marked the  
150th anniversary of Darwin's *On the  
Origin of Species*. How can we  
explain any genetic advantage for  
humans to carry so many microbes  
along? What benefit does our mi-  
crobiome provide us during health?  
And what happens during critical ill-  
ness? How does the intensive care  
unit (ICU) environment impact on our  
microbiota? What impact does SDD  
have on our microbiota? And what is  
the impact of SDD on the emergence  
of resistant bacteria in the ICU?

Zandstra et al. rightly allude to the  
fact that *Enterobacteriaceae* increase  
during critical illness—we also found  
a more than tenfold relative increase  
(as a percentage of total bacteria)

compared to data from healthy  
volunteers [4, 5]. To fight overgrowth  
of *Enterobacteriaceae* seems like a  
good idea—but is there a trade-off?  
Why did SDD—though providing a  
small but significant survival advan-  
tage compared to a control group—  
fail to further reduce mortality  
compared with patients receiving  
only oral topical antimicrobial  
products, even though these study  
participants had similarly increased  
percentages of *Enterobacteriaceae*?

A potential advantage of having  
such massive numbers of anaerobic  
bacteria in the colon is that colono-  
cytes feed on bacterial products.  
Indeed colonocytes feed on butyrate,  
produced by a limited number of  
colonic bacterial species. The *Fae-  
calibacterium prausnitzii* group is  
predominant amongst these butyrate-  
producing groups. This group is  
already reduced during tube feeding,  
resulting in a significantly reduced  
concentration of butyrate [4]. We  
describe a significant reduction in two  
groups of microbiota that help in  
maintaining the integrity of the large  
intestinal mucosa. Quoting Volleard's  
and Donskey's work, Zandstra et al.  
agree that SDD is a contradiction in  
terms. We apparently disagree in our  
concern that SDD might have a “dark  
side”, i.e. that some important bene-  
ficial microbiota are harmed by SDD.  
Besides, there is additional collateral  
damage: in the analysis of point-  
prevalence cultures from the de Smet  
study [6], an increase of intestinal  
colonisation by resistant organisms  
was observed after cessation of  
SDD [7].

None of the articles cited by  
Zandstra et al. describe the use of  
molecular methods for detection and/  
or enumeration of the intestinal mi-  
crobiota. Bacteria from the  
*F. prausnitzii* group are highly sensi-  
tive to oxygen and require very  
specific growth media. Culture-based  
quantification yields a high degree of  
culture bias. These limitations of

culture-based microbiological  
techniques have precluded reliable  
testing of the effects of antibiotics  
(i.e. SDD) on the intestinal micro-  
biota. We are obviously not the first to  
claim that SDD is not selective. We  
are, however, the first to back up  
these claims using absolute numbers  
of faecal bacteria derived from a  
quantitatively reliable, molecular  
method.

Referring to Wensinck as an  
argument that not *F. prausnitzii* but  
clostridia contribute to colonisation  
resistance (CR), Zandstra et al. dis-  
play a misunderstanding of the  
composition of the intestinal micro-  
biota. Although phenotypically  
Gram-negative, *F. prausnitzii* are  
genetically closely related to the  
Gram-positive clostridia Zandstra  
et al. refer to. In fact, they belong to  
clostridial cluster IV [8]. These phy-  
logenetic relations have only been  
discovered since the dawn of the  
molecular era.

We agree that the clinical impact  
of *F. prausnitzii* reduction in the  
critically ill is unclear—at least, that  
this should be further studied; we do  
not know whether there is a critical  
threshold of butyrate substrate for  
colonocytes to survive, or to maintain  
the integrity of the large intestinal  
mucosa. Recent studies have shown  
an anti-inflammatory effect of  
*F. prausnitzii* in Crohn's disease that  
could also be beneficial for the criti-  
cally ill [9]. We have, however,  
provided novel information that may  
explain the discrepancy of effective  
reduction of *Enterobacteriaceae* by  
SDD, which fails to translate into  
further survival benefit in comparison  
with selective oropharyngeal decon-  
tamination (SOD).

**Open Access** This article is distributed  
under the terms of the Creative Commons  
Attribution Noncommercial License which  
permits any noncommercial use, distribu-  
tion, and reproduction in any medium,  
provided the original author(s) and source  
are credited.

## References

1. Zandstra DF, Petros AJ, Silvestri L, de la Cal MA, Taylor N, Damjanovic V, van Saene HKF (2010) Selective decontamination of the digestive tract (SDD): selectivity is not required. *Intensive Care Med.* doi: [10.1007/s00134-010-1945-y](https://doi.org/10.1007/s00134-010-1945-y)
2. Benus RF, Harmsen HJ, Welling GW, Spanjersberg R, Zijlstra JG, Degener JE, van der Werf TS (2010) Impact of digestive and oropharyngeal decontamination on the intestinal microbiota in ICU patients. *Intensive Care Med.* doi: [10.1007/s00134-010-1826-4](https://doi.org/10.1007/s00134-010-1826-4)
3. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, Nielsen T, Pons N, Levenez F, Yamada T, Mende DR, Li J, Xu J, Li S, Li D, Cao J, Wang B, Liang H, Zheng H, Xie Y, Tap J, Lepage P, Bertalan M, Batto JM, Hansen T, Le Paslier D, Linneberg A, Nielsen HB, Pelletier E, Renault P, Sicheritz-Ponten T, Turner K, Zhu H, Yu C, Li S, Jian M, Zhou Y, Li Y, Zhang X, Li S, Qin N, Yang H, Wang J, Brunak S, Doré J, Guarner F, Kristiansen K, Pedersen O, Parkhill J, Weissenbach J, MetaHIT Consortium, Bork P, Ehrlich SD, Wang J (2010) A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 464:59–65
4. Benus RF, van der Werf TS, Welling GW, Judd PA, Taylor MA, Harmsen HJ, Whelan K (2010) Association between *Faecalibacterium prausnitzii* and dietary fibre in colonic fermentation in healthy human subjects. *Br J Nutr* 29:1–8
5. Harmsen HJ, Raangs GC, He T, Degener JE, Welling GW (2002) Extensive set of 16S rRNA-based probes for detection of bacteria in human feces. *Appl Environ Microbiol* 68:2982–2990
6. de Smet AM, Kluytmans JA, Cooper BS, Mascini EM, Benus RF, van der Werf TS, van der Hoeven JG, Pickkers P, Bogaers-Hofman D, van der Meer NJ, Bernards AT, Kuijper EJ, Joore JC, Leverstein-van Hall MA, Bindels AJ, Jansz AR, Wesselink RM, de Jongh BM, Dennessen PJ, van Asselt GJ, te Velde LF, Frenay IH, Kaasjager K, Bosch FH, van Iterson M, Thijsen SF, Kluge GH, Pauw W, de Vries JW, Kaan JA, Arends JP, Aarts LP, Sturm PD, Harinck HI, Voss A, Uijtendaal EV, Blok HE, Thieme Groen ES, Pouw ME, Kalkman CJ, Bonten MJ (2009) Decontamination of the digestive tract and oropharynx in ICU patients. *N Engl J Med* 360:20–31
7. Oostdijk EA, de Smet AM, Blok HE, Thieme Groen ES, van Asselt GJ, Benus RF, Bernards SA, Frénay IH, Jansz AR, de Jongh BM, Kaan JA, Leverstein-van Hall MA, Mascini EM, Pauw W, Sturm PD, Thijsen SF, Kluytmans JA, Bonten MJ (2010) Ecological effects of selective decontamination on resistant gram-negative bacterial colonization. *Am J Respir Crit Care Med* 181:452–457
8. Duncan SH, Hold GL, Harmsen HJ, Stewart CS, Flint HJ (2002) Growth requirements and fermentation products of *Fusobacterium prausnitzii*, and a proposal to reclassify it as *Faecalibacterium prausnitzii* gen. nov., comb. nov. *Int J Syst Evol Microbiol* 52:2141–2146
9. Sokol H, Pigneur B, Watterlot L, Lakhdari O, Bermúdez-Humarán LG, Gratadoux JJ, Blugeon S, Bridonneau C, Furet JP, Corthier G, Grangette C, Vasquez N, Pochart P, Trugnan G, Thomas G, Blottière HM, Doré J, Marteau P, Seksik P, Langella P (2008) *Faecalibacterium prausnitzii* is an anti-inflammatory commensal bacterium identified by gut microbiota analysis of Crohn disease patients. *Proc Natl Acad Sci U S A* 105:16731–16736

R. F. Benus · H. J. Harmsen ·  
T. S. van der Werf (✉)  
University Medical Center Groningen,  
Internal Medicine, Hanzeplein 1,  
P.O. Box 30001, 9700 RB Groningen,  
The Netherlands  
e-mail: [t.s.van.der.werf@int.umcg.nl](mailto:t.s.van.der.werf@int.umcg.nl)  
Tel.: +31-50-3611501  
Fax: +31-50-3613216